

MEDICATIONS FOR WEIGHT MANAGEMENT

Medication	1.Saxenda	2.Contrave	3.Wegovy	4.Xenical
	(Liraglutide) 3.0mg	(Bupropion/Naltrexone)	(Semaglutide) 2.4mg	(Orlistat)
	DIN#02437899	DIN#02472945		DIN#02240325
What is it?	A hormone called GLP1.	A combination	A hormone called GLP1.	Gastrointestinal lipase
		medication		inhibitor.
		(antidepressant and		
		opioid antagonist).		
What does it	Creates a feeling of	Decreases appetite and	Creates a feeling of	Blocks the action of
do?	fullness, leads to	can help with cravings.	fullness, leads to	lipase and decreases
	improvement in portion		improvement in portion	the absorption of fat
	control.		control.	in the intestines.
Weight loss	Average of 8% below	Average of 8% below	Average of 15% below	Average of 3% below
expected	starting weight.	starting weight.	starting weight.	starting weight.
How is it	Daily injection, usually	2 pills twice daily (titrate	Weekly injection,	One pill three times
taken	in the evening.	up slowly).	usually in the evening.	with meals.
Side Effects	-Nausea (decreases	-Nausea (decreases	-Nausea (decreases	-Oily, loose bowel
	within weeks).	within weeks).	within weeks).	movements –
	-Gallstones and acute	Contraindication:	-Gallstones and acute	frequently leads to
	pancreatitis are rare.	Chronic opioid use,	pancreatitis are rare.	discontinuation.
		seizure disorder		
Cost	Approximately \$250-	Approximately	Approximately \$250-	Approximately
	\$500/month depending	\$325/month for full dose	\$500/month depending	\$180/month
	on dose.		on dose.	
	Martinize.			
	Saxenda'		Nordina-4504/16 to a scient	- and the second s
	liraglutide injection 6 mgmL/5x3 mL pref ed aar)		2.4 mg	Xenical [®] 120 mg
	Sol, For for metters prediction Viewtoree.sum (#)	120 tablets	wegovy [™] (semaglutide) injection	hard capsules Orlistat
	Sime is a singular of the U. How we have don't be used if any use is a final similar to above 30 Core on and "special" of 24 C do not of the set of models statistic for each of the set of the U. How the Special Similar for the set of models and the Subset.	(naltrexone HCl and bupropion HCl) Extended-Release Tablets	2.4 mg/0.75 mL twittinguy times weat 4 Single Doze Prefilled Pens	120 mg
	Keep the open moves of the test. Discard sen 31 days after first test.	8mg/90mg Attention: Dispense with the attached Medication Guide	Sadt gen derivers a single donr all 2.4 rag semaglatide Ter solersmen mei ervit linge Deer and	Oral use
	Baerar Grad	Tablets should not be cut, chewed, or crushed.	Backer Unigers year, Freder Literature Disponse The archivest Medication Sache to sach patient.	
	politicii	R only	non-meal	CID 84 hard capsules

How long do I have to use it? We strongly suggest that you start another effective method of weight loss maintenance if you stop the medication. This medication is not a quick fix or short-term solution. If you stop this medication, the weight that was lost can return.

This is for informational purposes only and has been prepared by the Wharton Medical Clinic -April 2022 www.whartonmedicalclinic.com